

## OBSERVABLE CONSEQUENCES OF MERGER-DRIVEN GAPS AND HOLES IN BLACK HOLE ACCRETION DISKS

KAYHAN GÜLTEKİN<sup>1</sup> AND JÖN M. MILLER<sup>1</sup><sup>1</sup>Department of Astronomy, University of Michigan, 500 Church Street, Ann Arbor, MI 48109, USA [kayhan@umich.edu](mailto:kayhan@umich.edu), [jonmm@umich.edu](mailto:jonmm@umich.edu).*Draft version July 3, 2012*

## ABSTRACT

We calculate the observable signature of a black hole accretion disk with a gap or hole created by a secondary black hole embedded in the disk. We find that for an interesting range of parameters of black hole masses ( $\sim 10^6$ – $10^9 M_\odot$ ), orbital separation ( $\sim 1$  AU to  $\sim 0.1$  pc), and gap width (1–19 disk scale heights), the missing thermal emission from a gap manifests itself in an observable decrement in the spectral energy distribution. We present observational diagnostics in terms of powerlaw forms that can be fit to line-free regions in AGN spectra or in fluxes from sequences of broad filters. Most interestingly, the change in slope in the broken powerlaw is almost entirely dependent on the width of gap in the accretion disk, which in turn is uniquely determined by mass ratio of the black holes, such that it scales roughly as  $q^{5/4}$ . Thus one can use spectral observations of the continuum of bright active galactic nuclei to infer not only the presence of a closely separated black hole binary but also the mass ratio. When the black hole merger opens a hole in the inner disk, the broad band SED of the AGN or quasar may serve as a diagnostic. Such sources should be especially luminous in optical bands but intrinsically faint in X-rays (i.e., not merely obscured). We briefly note that viable candidates may have already been identified.

*Subject headings:* black hole physics — galaxies: active

## 1. INTRODUCTION

The prevalence of supermassive black holes (SMBHs) at the centers of galaxies is well established (Richstone et al. 1998). In the context of a hierarchical merging universe, merging galaxies can lead to the merging of the black holes (BHs Volonteri et al. 2003). Mergers of BHs are important as one pathway for the growth of BHs and have even been suggested as the principal cause of the scaling relations between BH mass and host galaxy properties (Jahnke & Macciò 2011). Mergers are also strong gravitational wave emitters, and the asymmetric emission of gravitational waves, especially from spinning BHs, can lead to a recoil of the merged BH large enough to kick it out of the host galaxy potentially influencing the BH occupation fraction of galaxies (e.g., Volonteri et al. 2008; Baker et al. 2008; Merritt et al. 2004). Even if two galaxies with BHs merge and the BHs sink to the center of the merged galaxy, however, it is not a given that the BHs will merge within a Hubble time (e.g., Milosavljević & Merritt 2003).

The search for precursors to BH mergers has naturally focused on active galactic nuclei (AGNs) that are either spatially resolvable at separations of  $\sim 1$  kpc (e.g., Comerford et al. 2012; Blecha et al. 2012) or have spectroscopically distinct broad line regions at separations of  $\sim 0.1$  pc (e.g., Gaskell 1996; Boroson & Lauer 2009; Smith et al. 2010; Eracleous et al. 2011).

One avenue for finding BH pairs at very small separations comes from an analogy with protoplanetary disks in which the presence of a planet may be inferred from its influence on the protoplanetary disk, particularly through gaps and holes in the disk carved out by the planet (Goldreich & Tremaine 1980; Takeuchi et al. 1996; Armitage & Rice 2005; Espaillat et al. 2008). In this letter, we consider the observable consequences of an accretion disk gap caused by a secondary BH. Owing to the complexity of AGN and quasar spectra, we have developed observational diagnostics in terms of powerlaw forms that can be fit to line-free regions in spectra, or potentially

even to fluxes derived from sequences of broad filters. We have also given consideration to signatures that may be evident in broad-band SEDs of quasars and AGN that span the optical and X-ray regimes.

## 2. DESCRIPTION OF THE PHYSICAL SYSTEM

The presence of a (proto-)planet in a proto-planetary disk in orbit around a central star can lead to transfer of angular momentum such that the planet moves inwards (Type I migration; Goldreich & Tremaine 1980). Under certain circumstances (section 3), the planet can clear material near its radius, opening up a gap. Dynamical interactions between the planet and material at the gap edges tend to drive both the gap and the planet inward (type II migration; Takeuchi et al. 1996).

We are interested in the observational consequences of analogous systems for SMBHs. Consider a central BH of mass  $M$  with an accretion disk around it with a secondary BH of mass  $m$ , mass ratio  $q = m/M$ , in the plane of the disk at an orbital radius of  $a$ . If the secondary BH can open a gap in the accretion disk, the relative difference in emission would create an observational signature that could be used as evidence of close SMBH binary pairs at separations of  $\sim 10^{-3}$  pc.

We assume that the accretion disk is as described by Shakura & Sunyaev (1973), i.e., an  $\alpha$ -disk. The  $\alpha$ -disk model is necessarily a mathematically convenient approximation rather than a fully physical model, but it is still a successful model in many respects for a wide variety of astrophysical accretion systems. Our results are not bound to the Shakura & Sunyaev (1973) disk description, and we consider variations in section 5. What we require is that (i) azimuthally averaged disk midplane temperature is monotonic with radius in the inner  $\sim 10^3 R_G$ , where  $R_G = GMc^{-2}$  and (ii)  $h/r$ , the ratio of disk scale height to radius, is roughly constant. The recent considerations of azimuthal variations in disk temperature (Dexter & Agol 2011; Dexter & Quataert 2012) still assume that the scale of the variability does not change with

radius so that for a given variability model, each azimuthally averaged annular emission is still a function only of  $r$ . The secondary hole is assumed to have already found its way into the accretion disk of the primary and settled into a coplanar orbit as has been shown to occur (Bogdanović et al. 2007; Dotti et al. 2010). As we discuss below, the mass ratios of interest are  $q > 0.003$ . Below about  $q = 0.1$ – $0.01$ , a potential concern is that the mergers with halos containing small black holes will lead to a complete dispersal of the secondary halo before the BH can settle to the center. Recent numerical simulations, however show that it is possible for mergers with dwarf satellites can lead to secondary BHs of mass ratios  $q = 0.01$ – $0.001$  getting close enough to the primary that local processes can bring the BHs together (Bellovary et al. 2010). It is also possible that intermediate-mass BHs could form in situ in the accretion disk and would naturally be coplanar and have small mass ratios (McKernan et al. 2012).

### 3. CONDITIONS FOR GAP OPENING

There are two requirements that must be met in order for a gap in the disk to open: (1) the Hill sphere radius of the secondary must be larger than the disk scale height and (2) gap closing timescale from the viscous reaction of the disk is longer than the gap opening timescale.

#### 3.1. Hill Sphere Calculation

The Hill sphere defines the region around an orbiting body in which it is gravitationally dominant and has radius:

$$R_H \equiv 4 \left( \frac{m}{3M} \right)^{1/3} = 4 \left( \frac{q}{3} \right)^{1/3}. \quad (1)$$

The secondary BH will clear a gap in the accretion disk if  $R_H \geq h$ , which happens only when  $q \geq 3(h/r)^3$ . In the gas pressure dominated region where the binary orbits are most detectable,  $h/r = 0.004$ – $0.008$  for thin disks. We take a conservative value of  $h/r = 0.01$  as our fiducial value implies that  $q \geq 3 \times 10^{-6}$  and even minor mergers are sufficient to carve a gap of width  $w \sim 2R_H$ .

#### 3.2. Gap closing

Gaps will only exist if the gravitational torques open them faster than viscous diffusion can close them. The timescale for opening a gap

$$t_{\text{open}} \sim \frac{1}{m^2 q^2 \Omega} \left( \frac{w}{r} \right)^2, \quad (2)$$

where  $\Omega$  is the angular frequency of the secondary and  $m = r\Omega c_s^{-1}$  is the order of the Linblad resonance driving the gap creation (Takeuchi et al. 1996). The timescale to close the gap is

$$t_{\text{close}} \sim \frac{w^2}{\nu}, \quad (3)$$

where  $\nu$  is the kinematic viscosity. Requiring  $t_{\text{close}} > t_{\text{open}}$ , yields the requirement that

$$q > \left( \frac{c_s}{r\Omega} \right)^2 \alpha^{1/2} = \left( \frac{h}{r} \right)^2 \alpha^{1/2}, \quad (4)$$

where  $\Omega$ ,  $c_s$ , and  $h$  are all taken to be at the radius of the secondary. This requirement is equivalent to  $q > 10^{-5}$  for our fiducial disk parameters.

#### 3.3. Longevity of the gap

In order for a gapped accretion disk to be of any observational consequence, it needs to persist for a reasonably large fraction of an AGN's lifetime. If the angular momentum transferred from the secondary to the outer disk is greater than the angular momentum transferred from the inner disk to the secondary, then the secondary and the gap will both migrate inwards. The gap migration timescale when the local disk mass dominates over the secondary mass is (Takeuchi et al. 1996)

$$t_{\text{migr}} = \frac{2}{3\alpha} \left( \frac{h}{r} \right)^{-2} \Omega^{-1}. \quad (5)$$

For a  $10^8 M_\odot$  primary with a secondary at  $\sim 10^3 R_G$ , this corresponds to  $\sim 3 \times 10^5$  yr. For higher mass secondaries, the gap migration time is lengthened in the satellite-dominated regime by a factor of a couple over this value so that the lifetime of the gap is  $\sim 10^6$  yr.

When the BHs are close to each other, gravitational radiation is strong enough to shrink the orbit on interesting timescales. The Peters (1964) orbit-averaged evolution of the semimajor axis of two masses in orbit around each other with eccentricity  $e$  is

$$\frac{da}{dt} = -\frac{64G^3 Mm(M+m)}{5c^5 a^3 (1-e^2)^{7/2}} \left( 1 + \frac{73}{24}e^2 + \frac{37}{96}e^4 \right), \quad (6)$$

where  $G$  is the gravitational constant and  $c$  is the speed of light. For the case we consider,  $e = 0$ . The width of the gap is  $R_H$ , and the timescale for the secondary BH to move out of the gap is

$$T_g = R_H \left( \frac{da}{dt} \right)^{-1} = \frac{5c^5 a^4}{64G^3 Mm(M+m)} \left( \frac{m}{3M} \right)^{1/3}. \quad (7)$$

Taking  $a = r = 10^3 R_G$  and  $q = 10^{-2}$ , then  $T_g \approx 2 \times 10^7 (M/10^8 M_\odot)$  yr, much longer than other relevant timescales.

So for a gap lifetime of  $\sim 10^6$  yr and a typical AGN lifetime of  $\sim 10^7$  yr (Martini 2004), we can see  $10^{-1}$  of AGNs with companions with the right mass ratio and with separations  $r < 2 \times 10^3 R_G$ .

### 4. EVOLUTION OF THE GAP

Once a gap is established it will evolve along three possible routes when considering only viscous effects. At the beginning it is a gapped disk, effectively a standard disk with an annulus of width  $w$  missing. If the gap migration timescale is faster than or comparable to the timescale for accretion of the disk material inside of the gap, then the gap will have constant  $w/h$  with a decreasing radius.

If the mass accretion rate for the inner disk is faster than the gap migration rate, then it will partially decouple from the outer disk. This will first lead to an effective widening of the gap from the inner edge, making it more detectable. The inner disk will be continuously depleted and replaced by any material that accretes across the gap. If the mass accretion rate across the gap is the same as if there were no gap, then there would be no difference. On the other hand, if the accretion rate is a small fraction of what it would have been otherwise, the inner disk would be replaced by an accretion disk of a smaller accretion rate.

The details of how much accretion will persist across a gap created by a satellite are the subject of current numerical simulations. For the small mass ratios that are our primary interest here, however, it is plausible that the accretion rate is only reduced to 10% of what it would be in the absence of a satellite, i.e.,  $\dot{m}_{\text{red}} = 0.1\dot{m}$ , where  $\dot{m} \equiv \dot{M}/\dot{M}_{\text{Edd}}$  and  $\dot{M}_{\text{Edd}} \equiv L_{\text{Edd}}\eta^{-1}c^{-2} \approx 3 \times 10^{-8}(M/M_{\odot})M_{\odot} \text{ yr}^{-1}$  for an assumed  $\eta = 0.1$ .

For higher mass ratios, it is plausible that very little matter will accrete past the gap, resulting in essentially no replacement of the inner gap. In such a case, instead of a gap, there would be a hole in central accretion disk. The observational consequences of an accretion disk with a central hole—and the possible electromagnetic signal once the BHs merge—have been discussed before (Milosavljević & Phinney 2005; O’Neill et al. 2009), and we return to this in section 6.

The time to accrete the entire inner disk is

$$t_a = M_{\text{in}}\dot{M}^{-1}, \quad (8)$$

where  $\dot{M}$  is the mass accretion rate and  $M_{\text{in}}$  is the mass of the inner accretion disk. For our fiducial accretion disk, we take the BH to be accreting at a fraction  $\dot{m} = 0.01$ . The largest orbit we consider is roughly at the boundary outside of which free-free opacity dominates over Thomson scattering so that the mass inside the gap is roughly

$$M \approx 6 \times 10^{-11} \left( \frac{M}{M_{\odot}} \right)^{11/5} \alpha^{-4/5} \dot{m}^{23/15} M_{\odot} \quad (9)$$

so that the accretion timescale is

$$t_a \approx 0.002 \alpha^{-4/5} \left( \frac{M}{M_{\odot}} \right) \dot{m}^{6/5} \dot{m}^{8/15}. \quad (10)$$

For our fiducial case,  $t_a \approx 3 \times 10^7 \text{ yr}$ , so that the inner disk is depleting itself more slowly than the gap’s inward migration. The above is only valid for  $q \ll 1$ . For mass ratios near unity, the dynamical torques will clear a hole inside of the orbit on a short timescale.

## 5. SPECTRAL ENERGY DISTRIBUTION OF A GAPPED ACCRETION DISK

The spectral energy distribution (SED) of a thermally emitting disk with temperature a function only of radius is

$$F_{\lambda}(\lambda) = \int_{R_{\text{in}}}^{R_{\text{out}}} B_{\lambda}[T(r')] g(r') 2\pi r' dr', \quad (11)$$

where  $B_{\lambda}(T)$  is the Planck function and  $g(r')$  is a function that describes the emissivity as a function of radius. We assume that  $g(r') = 1$  (optically thick) where the disk material is present (i.e.,  $R_{\text{in}} < r' < r - w$  and  $r + w < r' < R_{\text{out}}$ ) and  $g(r) = 0$  elsewhere.

The absence of a ring in a thermally emitting disk will show up in the SED as a broad, shallow dip. Without a gap, the SED will fall off as  $\lambda F_{\lambda} \sim \lambda^{-4/3}$ . With a gap, the SED will be appear roughly as a broken powerlaw with a slope that is steeper and shallower than  $\lambda^{-4/3}$  blueward and redward of the maximum deviation from the standard SED. While it is obviously possible to model an observed SED with the proper Planck function treatment, we use a broken powerlaw model to demonstrate the precision required to identify a gapped disk. We generated SEDs (Fig. 1) for gapped disks using equation (11) for various values of  $w$ ,  $r/R_G$ ,  $M$ , and  $\dot{m}$ . For

each simulated SED, we fitted broken powerlaw in the vicinity of the break of the form

$$\lambda F_{\lambda} = \begin{cases} \lambda^{-x} & \lambda \leq \lambda_0 \\ \lambda^{-y} & \lambda > \lambda_0 \end{cases}, \quad (12)$$

where  $x > 4/3$  and  $y < 4/3$  with an arbitrary normalization implied. The change in slope,  $\Delta \equiv x - y$ , from one side of the break to the other at  $\lambda = \lambda_0$  increases as the gap width increases. We find

$$\lambda_0 \approx 140 \left( \frac{r}{R_G} \right)^{3/4} \left( \frac{M}{10^8 M_{\odot}} \right)^{1/4} \left( \frac{\dot{m}}{0.01} \right)^{-1/4} f(w/h) \text{ \AA}, \quad (13)$$

where  $f(x)$  is a residual function that is smoothly varying over  $w/h$  and has absolute value  $|f(x) - 1| < 0.35$  for most of the parameter space of interest. The dependence on  $r$ ,  $\dot{m}$ , and  $M$  is easily understood since the peak wavelength of a blackbody scales as  $\lambda \sim T^{-1}$  and temperature and mass scale as  $T \sim \dot{m}^{1/4} r^{-3/4}$  in the outer regions where the secondary spends most of the time and  $M \sim T^{-1/4}$ . The dependence on  $w$  comes from the area weighting of the output SED. The change in slope is well approximated by

$$\Delta \approx 0.002(w/h)^{5/4} \quad (14)$$

for  $w/h < 160$ . Above  $w/h > 160$ , the change in slope is larger than the above approximation because the emitting annulus just outside of the gap is further down the Wien exponential drop off. Since  $w/h$  is purely a function of  $q$ , it is possible to infer the mass ratio of the binary from just the change in slope, independent of  $\dot{m}$ ,  $M$ , and  $r$ .

Rafikov (2012) found that a gapped accretion disk would have an SED such that  $\lambda F_{\lambda} \propto \lambda^{12/7}$ . If  $T(r) \sim r^{-0.9}$ , then the peak of the SED scales as  $\lambda F_{\lambda} \propto \lambda^{-1.7}$ , approximately as  $\lambda^{12/7}$ . There is no qualitative difference in the approach to modeling SEDs of such gapped disks. We find that the change in powerlaw slope for such a temperature power-law is only decreased by about 10% compared to our fiducial temperature profile. Thus one could use the overall slope of the continuum powerlaw to identify gapped accretion disks and then use detailed measurements of  $\Delta$  and  $\lambda_0$  to infer properties of the BHs.

## 6. SED OF AN ACCRETION DISK WITH A HOLE

For higher mass ratios and suitably small separations, almost the entire region inside of the orbit will be dynamically unstable as can be seen from considering the size of the Roche lobes for  $q \approx 1$  binaries. Thus all gas will be cleared out of this region, leaving an accretion disk with a hole of size  $r \sim 2a$  in the center. The effect of this on the SED is much more dramatic than that of a gap. Most or all of the shortest wavelength emission will be cut out, leaving a red SED.

The absence of an inner accretion disk owing to a BH merger event may therefore be identified through the broad-band SED. The outer accretion disk will still have a high accretion rate, and should be luminous in optical bands. Indeed, the outer disk may be abnormally luminous in optical, or even in UV bands (see, e.g., Kocsis et al. 2012), depending on the mass accretion rate and the truncation radius. On simple grounds, UV luminosity is not a good diagnostic: a holed disk around a BH of lower mass is observationally degenerate with a filled inner disk around a BH of higher mass. Rather,




**Figure 1.** All panels show simulated SEDs with gaps, with our fiducial parameters of  $M = 10^8 M_\odot$ ,  $h/r = 0.01$ , and assuming our standard temperature profile. The left and center panels assume  $r = 1000 R_G$  and show the effect of increasing the gap width with values  $w/h = 0, 10, 60, 80, 100, 140$ , and  $180$ . The central panel displays a zoomed in version of the same and shows the depth of the dip in the SED increasing as well as the differences in slopes on either side of the local minimum increase as  $w/h$  increases. Empirically, we find that the change in slope scales approximately as  $(w/h)^{5/4}$ , allowing one to infer the mass ratio of the binary from just the change in slopes. The right panel assumes  $w/h = 180$  and shows the effect of increasing the gap radius with values from  $r/R_G = 200$  to  $2000$  in steps of  $200$ . For reference, also plotted is a gapless SED. Modeled as a broken powerlaw near the gap, we find that the break occurs at a wavelength that scales with gap radius as  $\lambda \sim (r/R_G)^{3/4}$ . This comes from the temperature profile's dependence of  $T \sim r^{-3/4}$  and having the peak wavelength of a blackbody scale as  $\lambda \sim T^{-1}$ . The break wavelength also depends on  $M$  and  $\dot{m}$  as shown in equation (13).

diminished X-ray emission is a better diagnostic, since non-thermal X-ray emission is thought to tap into the deep gravitational potential close to the BH. Residual accretion within the hole may persist at a low level, but even this gives rise to X-ray variability in Sgr A\*. Thus, X-ray variability may be an additional hallmark. Thus, a source that is luminous in optical bands but intrinsically faint (not faint due to obscuration) and potentially variable in X-rays might signal a recent or pending BH merger event.

It is possible that such sources have already been identified. When the distance to a given AGN is known a priori, e.g., through spectroscopy, SDSS colors are able to distinguish whether or not a source is intrinsically luminous in, e.g., red bands, or just apparently red through obscuration. In particular,  $g - i$  is very effective, or  $\Delta(g - i)$ , where one measures color relative to the mean quasar color in a given red-shift bin. Hall et al. (2006) employed this technique to 3814 SDSS quasars with the aim of isolating obscured BHs. Follow-up observations with *Chandra* revealed that a subset of *intrinsically* red quasars are *not* X-ray-faint because of obscuration. In particular, seven sources show no evidence of even moderate intrinsic absorption in X-rays, and two are constrained to have internal column densities below  $2 \times 10^{21} \text{ cm}^{-2}$  (Hall et al. 2006). This suggests that such sources are rare ( $7/3814 = 1.8 \times 10^{-3}$ ), consistent with expectations based on the peculiar configuration required and the relevant timescales, and also consistent with expectations based on broader surveys (Gibson et al. 2008). Another interesting case for a quasar with a small inner hole in its accretion disk may be PHL 1811, which is optically very luminous but X-ray-faint (Leighly et al. 2004; Choi et al. 2005).

## 7. DISCUSSION AND CONCLUSIONS

We have described the basic observational appearance of a BH with a gapped or holed accretion disk, both of which are observable through their broad-band SEDs. We have focused our consideration on parameter space that results in observable signatures between  $2000 \text{ \AA}$  and  $2 \mu\text{m}$ . At shorter wavelengths, absorption of ultraviolet photons will prevent unambiguous determination of details of the AGN continuum. At longer wavelengths, reprocessed emission from dust grains will similarly contaminate thermal emission from an accretion disk. In this range, however, it is possible to measure the continuum emission to the precision needed to infer the pres-

ence of a gap. Current and future large surveys such as SDSS, Pan-STARRS, and LSST may be able to exploit the observational signatures we have developed.

This treatment has ignored some potential complications that will need to be considered in the future. Of chief concern is that we have treated the emission as only coming from the top of the disk. Emission coming from the walls of the gap will complicate the signal and, depending on the vertical temperature structure, either increase or decrease the observability of the gap. Tidal features produced by the secondary as seen in simulations of protoplanetary disks will also need to be considered and may lead to additional corroborating observational signals. The periods of the secondaries that we are observable are months to several years, allowing for a potential detection of a modulating signal.

We thank Cole Miller for extremely useful discussions. KG acknowledges support provided by NASA through Chandra Awards GO0-11151X and G02-13111X and through Hubble Award HST-GO-12557.01-A awarded by the STScI.

## REFERENCES

- Armitage, P. J., & Rice, W. K. M. 2005, preprint (arXiv:astro-ph/0507492)
- Baker, J. G., Boggs, W. D., Centrella, J., Kelly, B. J., McWilliams, S. T., Miller, M. C., & van Meter, J. R. 2008, *ApJ*, 682, L29
- Bellovary, J. M., Governato, F., Quinn, T. R., Wadsley, J., Shen, S., & Volonteri, M. 2010, *ApJ*, 721, L148
- Blecha, L., Loeb, A., & Narayan, R. 2012, preprint (1201.1904)
- Bogdanović, T., Reynolds, C. S., & Miller, M. C. 2007, *ApJ*, 661, L147
- Borison, T. A., & Lauer, T. R. 2009, *Nature*, 458, 53
- Choi, J., Leighly, K. M., & Matsumoto, C. 2005, in *Bulletin of the American Astronomical Society* 37, American Astronomical Society Meeting Abstracts #206, 447
- Cormier, J. M., Gerke, B. F., Stern, D., Cooper, M. C., Weiner, B. J., Newman, J. A., Madsen, K., & Barrows, R. S. 2012, *ApJ*, 753, 42
- Dexter, J., & Agol, E. 2011, *ApJ*, 727, L24
- Dexter, J., & Quataert, E. 2012, preprint (1206.0739)
- Dotti, M., Volonteri, M., Perego, A., Colpi, M., Ruszkowski, M., & Haardt, F. 2010, *MNRAS*, 402, 682
- Eracleous, M., Borison, T. A., Halpern, J. P., & Liu, J. 2011, preprint (1106.2952)
- Español, C., Calvet, N., Luhman, K. L., Muzerolle, J., & D'Alessio, P. 2008, *ApJ*, 682, L125
- Gaskell, C. M. 1996, *ApJ*, 464, L107
- Gibson, R. R., Brandt, W. N., & Schneider, D. P. 2008, *ApJ*, 685, 773
- Goldreich, P., & Tremaine, S. 1980, *ApJ*, 241, 425
- Hall, P. B., Gallagher, S. C., Richards, G. T., Alexander, D. M., Anderson, S. F., Bauer, F., Brandt, W. N., & Schneider, D. P. 2006, *AJ*, 132, 1977

- Jahnke, K., & Macciò, A. V. 2011, *ApJ*, 734, 92
- Kocsis, B., Haiman, Z., & Loeb, A. 2012, preprint (1205.5268)
- Leighly, K., Halpern, J., & Jenkins, E. 2004, in *Astronomical Society of the Pacific Conference Series* 311, AGN Physics with the Sloan Digital Sky Survey, ed. G. T. Richards & P. B. Hall, 277
- Martini, P. 2004, *Coevolution of Black Holes and Galaxies*, 169
- McKernan, B., Ford, K. E. S., Lyra, W., & Perets, H. B. 2012, preprint (1206.2309)
- Merritt, D., Milosavljević, M., Favata, M., Hughes, S. A., & Holz, D. E. 2004, *ApJ*, 607, L9
- Milosavljević, M., & Merritt, D. 2003, *ApJ*, 596, 860
- Milosavljević, M., & Phinney, E. S. 2005, *ApJ*, 622, L93
- O'Neill, S. M., Miller, M. C., Bogdanović, T., Reynolds, C. S., & Schnittman, J. D. 2009, *ApJ*, 700, 859
- Peters, P. C. 1964, *Physical Review*, 136, 1224
- Rafikov, R. R. 2012, preprint (1205.5017)
- Richstone, D., Ajhar, E. A., Bender, R., et al. 1998, *Nature*, 395, A14
- Shakura, N. I., & Sunyaev, R. A. 1973, *A&A*, 24, 337
- Smith, K. L., Shields, G. A., Bonning, E. W., McMullen, C. C., Rosario, D. J., & Salviander, S. 2010, *ApJ*, 716, 866
- Takeuchi, T., Miyama, S. M., & Lin, D. N. C. 1996, *ApJ*, 460, 832
- Volonteri, M., Haardt, F., & Gültekin, K. 2008, *MNRAS*, 384, 1387
- Volonteri, M., Haardt, F., & Madau, P. 2003, *ApJ*, 582, 559